

12 児童生徒の疾病異常（令和元年度・公立）

（学校安全保健課調）

| 区 分 | | 小学校 | | 中学校 | | 義務教育学校 （前期） | | 義務教育学校 （後期） | | 高等学校 （全日制） | |
|--|-------------------|-------|-------|-------|-------|----------------|-------|----------------|-------|---------------|-------|
| | | 男 | 女 | 男 | 女 | 男 | 女 | 男 | 女 | 男 | 女 |
| 裸 眼 視 力 | 1.0未満0.7以上の者 | 10.3% | 11.9% | 12.3% | 13.0% | 6.0% | 7.9% | 9.0% | 12.3% | 14.7% | 15.5% |
| | 0.7未満0.3以上の者 | 9.9% | 12.4% | 15.1% | 16.8% | 12.4% | 16.3% | 12.3% | 13.9% | 17.5% | 17.9% |
| | 0.3未満の者 | 4.8% | 6.2% | 11.5% | 14.7% | 2.3% | 5.6% | 9.0% | 4.1% | 16.3% | 16.4% |
| 眼 | 感染性眼疾患の者 | 0.1% | 0.1% | 0.0% | 0.0% | 0% | 0% | 0% | 0% | 0.1% | 0.1% |
| | その他の眼疾患・異常の者 | 6.9% | 6.0% | 6.7% | 5.8% | 6.9% | 4.6% | 7.3% | 3.1% | 3.5% | 3.2% |
| 聴 | 難聴の者 | 0.6% | 0.7% | 0.3% | 0.3% | 0% | 0% | 0% | 0% | 0.9% | 0.8% |
| 耳 ・ 鼻 ・ 咽 | 耳疾患の者 | 6.0% | 5.9% | 5.1% | 3.6% | 7.9% | 6.2% | 3.4% | 4.7% | 2.0% | 1.2% |
| | 鼻・副鼻腔疾患の者 | 13.4% | 9.3% | 12.2% | 9.0% | 39.5% | 42.3% | 20.7% | 17.1% | 8.0% | 6.8% |
| | 口腔咽頭疾患の者 | 1.2% | 0.9% | 0.4% | 0.3% | 0.7% | 0.8% | 0.9% | 0.8% | 0.3% | 0.2% |
| 歯 | 処置完了の者 | 21.1% | 19.5% | 17.5% | 20.2% | 23.9% | 22.8% | 24.1% | 25.1% | 23.8% | 28.6% |
| | 未処置歯のある者 | 21.0% | 19.1% | 14.0% | 14.6% | 27.0% | 32.1% | 27.0% | 35.7% | 16.4% | 16.2% |
| | その他の疾病の者 | 8.2% | 8.3% | 6.2% | 4.9% | 12.4% | 8.2% | 12.8% | 10.5% | 1.5% | 2.3% |
| 尿 | 蛋白陽性の者 | 0.4% | 1.0% | 1.8% | 2.2% | 1.3% | 0% | 0.7% | 0.6% | 1.3% | 1.5% |
| | 潜血陽性の者 | 0.4% | 1.1% | 0.6% | 3.3% | 0% | 0.5% | 1.3% | 2.3% | 0.5% | 2.5% |
| | 糖陽性の者 | 0.0% | 0.0% | 0.1% | 0.2% | 0% | 0% | 0% | 0.6% | 0.1% | 0.2% |
| 寄生虫 | 寄生虫卵保有の者 | 0.1% | 0.1% | - | - | 0% | 0% | - | - | - | - |
| そ の 他 の 疾 病 及 び 異 常 | 栄養不良の者 | 0.0% | 0.0% | 0.0% | 0.1% | 0% | 0% | 0% | 0% | 0.1% | 0.1% |
| | 肥満傾向の者 | 1.3% | 0.7% | 0.9% | 0.4% | 2.2% | 2.2% | 1.4% | 0.6% | 0.4% | 0.2% |
| | 脊柱側わんの者 | 0.4% | 0.9% | 1.0% | 4.7% | 1.8% | 1.1% | 7.4% | 7.4% | 0.9% | 2.7% |
| | その他の脊柱・胸郭・四肢の異常の者 | 0.7% | 0.5% | 1.2% | 0.9% | 2.7% | 2.2% | 0% | 0% | 1.0% | 0.7% |
| | 感染性皮膚疾患の者 | 0.1% | 0.1% | 0.0% | 0.0% | 0.4% | 0% | 0% | 0% | 0% | 0% |
| | アトピー性皮膚炎の者 | 4.4% | 3.9% | 4.4% | 3.8% | 6.2% | 7.1% | 7.4% | 4.5% | 2.4% | 1.9% |
| | 心臓疾患・異常の者 | 1.0% | 0.9% | 1.2% | 1.1% | 0% | 1.6% | 0.7% | 2.3% | 0.7% | 0.7% |
| | ぜん息の者 | 6.9% | 4.9% | 5.7% | 3.7% | 17.3% | 13.6% | 6.1% | 3.4% | 2.4% | 1.9% |
| 腎臓疾患の者 | 0.3% | 0.3% | 0.4% | 0.5% | 0.4% | 0.5% | 0.7% | 1.1% | 0.2% | 0.2% | |
| 結 核 | 精密検査を受けた者 | 0.4% | 0.2% | 0.1% | 0.1% | 0% | 0.5% | 0.6% | 0% | 1.1% | 1.0% |
| | 結核の者 | 0.0% | 0% | 0% | 0% | 0% | 0% | 0% | 0% | 0% | 0% |

注・定期健康診断結果による。

- ・「0」は皆無、「0.0」は数値が表章単位未満、「-」は実施しない場合である。