

マイワシの粗脂肪量

水揚日: 令和2年2月20日


船名: 38 たいよう丸 漁場: 36°12'N 140°48'E (鹿島沖)

Lipid content of Japanese sardines

Landing day : February 20, 2020

Fishing ground : Off the coast of Kashima, Japan

36°12'N 140°48'E



No.	体長cm Standard length	体重g Body weight	粗脂肪量% Lipid content	No.	体長cm Standard length	体重g Body weight	粗脂肪量% Lipid content
1	18.5	76.7	8.4	11	13.5	29.6	12.1
2	14.9	39.4	13.4	12	13.7	29.2	10.9
3	15.3	42.6	16.4	13	13.2	28.5	10.9
4	14.6	36.2	5.3	14	14.1	32.9	7.8
5	14.7	35.2	2.9	15	14.3	31.8	5.9
6	15.0	36.9	7.8	16	14.1	29.9	3.5
7	16.1	43.4	5.9	17	14.7	36.3	10.9
8	14.9	34.9	3.5	18	14.4	35.6	12.1
9	15.3	41.4	7.2	19	14.2	32.2	8.4
10	14.8	36.4	5.9	20	14.3	33.8	9.6

千葉県水産総合研究センター銚子分室 TEL 0479-24-9796

Chiba Prefectural Fisheries Reserch Center

千葉県農林水産技術会議